FEDERAL INSTITUTE OF INDUSTRIAL PROPERTY		
Selected databases Query		Abstract
parameters Query definition Refine query (Query results Basket Saved queries Statistics Help Proposals Exit	Status	of 28.03.2007 - operates
	(11) Number of the patent document	2214241
	(13) Kind of document	C2
	(14) Document date	2003.10.20 Search
	(19) Publishing country or organization	RU
	(21) Application number	9812357%/14
	(22) Application filing date	1997.06.23
	(24) Date started of validity of the patent	1997.06.23
	(31) Priority application number	60/022,004
	(32) Date of filing of priority application	1996.06.28
	(33) Alloting country or organization	US
	(43) Unexamined printed documents without grant	2000.09.27 Search
	(45) Date	2003.10.20 Search
	(516) Edition of IPC	7
	(51) Main classification IPC	A61K31/35 Search IPC
	(51) Main classification IPC	A61P3/04 Search IPC
	Title	METHOD FOR TREATMENT OF OBESITY AND METHOD FOR INDUCING WEIGHT LOSS IN MAMMALS
	(71) Applicant information	ORTO-MAKNEJL FARMAS'JuTIKAL, INK. (US) Search
	(72) Inventor information	ShEhNK Richard P. (US) Search
	(73) Grantee (asignee) information	ORTO-MAKNEJL FARMAS'JuTIKAL, INK. (US) Search
	(74) Attorney, agent, representative information	Lebedeva Natal'ja Georgievna
	(85) PCT date art. 22/39	1998.12.25
	(86) PCT or regional filing information	US 97/10953 (23.06.1997)
	(87) PCT or regional filing information (publ.)	WO 98/00130 (08.01.1998)
	(98) Mail address	129010, Moskva, ul. B. Spasskaja, 25, str.3, OOO

"Juridicheskaja firma Gorodisskij i Partnery", pat.pov. N.G.Lebedevoj

Abstract

## FEDERAL INSTITUTE OF INDUSTRIAL PROPERTY

Selected databases
Query parameters
Query definition
Refine query
Query results
Basket
Saved queries
Statistics
Help

roposals Exit Bibliography #2214241. Abstract

FIELD: medicine therapy endocrinology SUBSTANCE: inve

FIELD: medicine, therapy, endocrinology. SUBSTANCE: invention relates to treatment of obesity. Invention proposes to use topiramate in daily dose 25-400 mg. Method provides significant reduction of weight gain rate in mammals due to diminishing metabolic effectiveness of food, in part, with selective reduction of formation and accumulation of lipids in adipose tissue. EFFECT enhanced effectiveness of treatment method. 3 cl, 3 tbl, 1 ex

Bibliography

